

## Nunavut Routine Adult Immunization Schedule

Vaccine	Age Group		
	19 – 24 Years	25 – 49 Years	≥ 50 Years
Influenza 1	1 dose recommended yearly.		
Meningococcal-	See guidelines		
C-ACYW <sup>2</sup>	below		
Tetanus,	1 booster dose recommended every 10 years throughout adulthood.		
Diphtheria (Td) <sup>3</sup>			
Varicella 4	See guidelines below and review Varicella Immunization Protocol		
Pneumococcal			See guidelines below
Polysaccharide			
(Pneu-P-23) <sup>5</sup>			
Hepatitis B	3 doses series recommended for select high-risk adults		
Vaccine 6			

- 1. Follow annual influenza guidelines.
- If not previously vaccinated with meningococcal vaccine in adolescence (including previous vaccination with Meningococcal C vaccine) 1 dose is recommended for students or athletes <25 years of age. Review immunization protocol for high-risk eligibility.
- 3. 1 dose in adulthood should include Acellular Pertussis (Tdap).
- 4. 2 doses at least 6 weeks apart. Recommended for susceptible (non-immune) health care workers and women who test negative for varicella antibodies at prenatal screen (to be given in postpartum period).
- 5. 1 dose recommended to all adults in this age group. 1 additional booster dose recommended 5 years later for select high-risk adults. Refer to Pneumococcal Polysaccharide 23-valent (Pneu-P-23) immunization protocol for high-risk eligibility. Maximum number of doses is 2.
- 6. 3 dose series recommended for select high-risk adults. Refer to Hepatitis B Immunization Protocol for eligibility and dose series guidelines.