



DELIMITING NUNAVUT CARIBOU POPULATIONS USING NUCLEAR DNA

Mitch Campbell¹

¹Regional Wildlife Biologist, Department of Environment, Government of Nunavut,
P.O. Box 120, Arviat, NU X0C 0E0

November 2006

Status Report, No. 24

Campbell, M. 2006 (Nov.). Delimiting Nunavut caribou populations using nuclear DNA. Government of Nunavut, Department of Environment, Status report: 24, Iqaluit, 8 pp.

**INTERIM REPORT
TO THE NUNAVUT WILDLIFE MANAGEMENT BOARD**

Prepared by: The Department of Environment
Kivalliq Region
P.O. Box 120
Arviat, NU. X0C 0E0

November, 2006

Title: Delimiting Nunavut Caribou Populations Using Nuclear DNA.

Project Leader: Mitch Campbell
Regional Wildlife Biologist
Department of Environment
Box 120
Arviat, NU.
X0C 0E0
Ph: (867)857-2828
Fax: (867)857-2828
e-mail: mcampbell@gov.nu.ca

Summary:

The management of Nunavut caribou and muskox populations as distinct demographic units with associated harvesting recommendations requires research methodologies capable of delimiting these populations. Studies of caribou and muskox movement, population trends, and seasonal range will be used to direct the proposed study as well as compliment its results. Many methods have the potential of meeting these research goals however a genetic approach is proposed as a cost effective general first step towards defining demographic units to caribou and muskox populations across Nunavut.

Background:

Caribou (*Rangifer tarandus*) and Muskox (*Ovibos moschatus*) are valuable economic and cultural game species in Nunavut. Through traditional knowledge and scientific studies we know that these species, especially caribou, are not always a dependable resource due to the unpredictable nature of range shifts, population declines and resource availability. In the case of Peary caribou a changing environment has led to declines in many populations leading to their addition to the COSEWIC endangered species list. In addition to natural events

many caribou and muskox populations have and will experience various levels of stress due to human impacts on their ranges. As caribou and muskox range conditions change with increased land use, the maintenance of existing genetic diversity could become more difficult ((Zittlau, 2004). Genetic diversity is considered necessary if a population is to avoid the risk of inbreeding effects and adapt to changing environmental conditions (Zittlau, 2004; Proctor, 2003; Proctor and Paetkau, 2004).

It is important to understand the genetic structure of caribou populations and to determine the current levels of diversity within these populations (Zittlau, 2004; Proctor, 2003; Proctor and Paetkau, 2004). The importance of this kind of study reaches into the very management regime and decision-making process where management decisions effecting one herd can have profound effects on related herds (Zittlau, 2004).

Objectives:

Much debate has surfaced over the years concerning the relatedness of caribou populations and muskox distributions across northern North America. Specific to this report is the need to delimit these populations and or groupings for many reasons:

- 1) To add depth to Nunavuts current understanding of caribou and muskox populations and management zones. Biological rationale:
- 2) To help wildlife managers understand the conservation issues involved when faced with local removals of caribou and/or muskox through either harvesting or natural events. Biological rationale:
- 3) To gain insight into the degree of mixing that may occur between populations. Biological rationale:.
- 4) To help define management units and associated harvesting recommendations such as total allowable harvest (TAH) etc to insure both conservation for and availability and accessibility to Nunavumiut. Biological rationale:.

Application of Results:

The proposed research is meant to provide a general knowledge base to both complement and help focus more specific population research including both future as well as ongoing telemetry studies, focused genetic studies, and demographic studies. The results of the study will be used to apply a more demographic approach to the management and conservation of Nunavut caribou

and muskox populations. This research will also help to build the knowledge base necessary for a more predictive management approach. The results will also be used to direct future research as well as complement existing population and range studies pulling together both past and present research into a more demographic approach to the conservation of Nunavut wildlife.

Study Area:

The study area takes in the entire Nunavut territory and will include barren ground caribou (*Rangifer tarandus groenlandicus*), peary caribou (*Ranger tarandus peari*) and muskox (*Ovibos moschatus*) populations and/or evidenced aggregations. (Figure 1).

Project Design:

The intent is to provide sufficiently continuous coverage to allow detection and quantification of genetic subdivision. The strategy varies between species because of differences in population organization. For caribou we intend to sample 30 to 50 samples from each recognized herd of caribou. Muskoxen are distributed more continuously in space, and the option exists to obtain continuous sample coverage or to collect clusters of samples at regular intervals. The latter approach will be used because it fits with opportunistic collection exemplified through traditional harvesting methods. This approach also allows us to make 'point' estimates of within-population genetic variation.

Each sample from either species will be analyzed using 12 highly variable microsatellite markers. Markers will be selected to allow maximum overlap with published data, particularly for caribou where eight-locus microsatellite data have been collected for many herds (thesis chapters 3 & 4). The allele scoring done at WGI has been calibrated relative to the scoring used by Zittlau (2004), making it possible to merge any new data with existing results.

Once the microsatellite data have been generated, they can be analyzed — and re-analyzed at any time in the future with any number of programs that attempt to describe how genetic variation is partitioned within and between populations, and what that partitioning indicates about connectivity between populations. One of our core analyses would be to use the Bayesian clustering program Structure 2.1 (Pritchard, 2000). This program partitions samples into genetic clusters that are defined without reference to geographic location. Zittlau (2004) failed to identify distinct subgroups of Barren Ground caribou with this approach, and one of our goals would be to test whether increasing sample sizes, both in terms number of samples and number of locations in the genome (markers), would uncover population subdivisions that we didn't have the power to detect with existing data. Most other population genetics programs refer to user-defined population groups

populations where only historical samples exist other tissue samples can be used as deemed appropriate by the genetic lab. Where possible whole right ears or the tips of tails were collected to insure the sampling of only one animal. Geographic location, date, estimated age, sex, group size and the hunters name accompanied each sample in a sample bag provided. Up to 100 samples were requested from each community wildlife office at an average cost of five dollars per sample going to the hunter. Immediately following the sampling period samples were and continue to be shipped to the Arviat office for cataloging and final shipment to a DNA profiling facility for analysis and reporting. The expected completion for analysis is March 2007 and for reporting August 2007.

Results:

Muskox tissue sampling and analysis 2005 to 2007

Sample collection was initiated April 1st, 2005 and is ongoing across Nunavut. Delays in the collection of samples have set the project back one year. To date the majority of the Nunavut wide muskox tissue samples have been collected from each of four locations; Kugluktok, Arviat, Baker Lake and Grise Fiord. The completion of the muskox sampling is scheduled for late February 2007. The preparation of these samples for shipment and lab analysis is scheduled for completion by March 2007.

Caribou tissue sampling and analysis 2005 to 2007

Collection of caribou samples was initiated April 1st, 2005 and is ongoing across Nunavut. To date samples from the Qamanirjuaq, Southampton Island, Ellsmere Island and the Northeast Mainland populations of caribou have been collected. Collections for the remaining herds are on going and expected to be complete by late February 2007. Sample collections from the identified Peary caribou populations have been problematic due to the low densities and extremely low harvest rates. Efforts are currently being made to locate archived genetic material to meet study protocols. Completion of genetic analysis and final reporting is scheduled for early march and late August 2007 respectively.

Management Implications:

The goals and objectives of this study are to examine the demography of Nunavuts caribou populations through the examination of genetic relatedness. This information can then be used with other demographic studies in an attempt to define caribou populations as distinct populations for the purposes of harvest management and future conservation. An additional goal of the Nunavut Wildlife Research Group will be the development of population based management plans into which the results of this and other studies will be fed.

Community Consulting and Reporting:

Formal consultations regarding this project have been completed within seven Kivalliq communities and have been discussed opportunistically at regular HTO meetings Nunavut wide. Broad support for the project has been received throughout Nunavut. Proposed reporting of the results of this research will take place following the genetic analysis on or about August 2007.

Schedule:

Project: Delimiting Nunavut Caribou Populations Using Nuclear DNA.				
Output or Step	Start Date	End Date	Person Days	Overtime Est. (hrs)
Community meetings (Update)	Nov (2006)	Nov (2006)	9	
Sampling - All Kivalliq	April(2005)	Feb. (2007)	7	
Sampling – Baffin	April(2005)	Feb. (2007)	7	
Sampling - Kitikmeot	April(2005)	Feb. (2007)	7	
Analysis	April (2006)	March (2007)		
Study Completion (All phases)	Feb. (2007)	Aug. (2007)		
Totals			30	0

Anticipated Outputs:

- 1) Interim and final reports to DoE, contributing agencies, community HTOs and co-managers.
- 2) Distribution of the Kivalliq Wildlife Research Update twice a year (Oct. and Mar.)
- 3) DoE file report following study completion.
- 4) Scientific papers in refereed journals following the completion of the project.
- 5) Community meetings and/or radio show describing methods and benefits of research and to obtain recommendations and/or address concerns.

Personnel:

- Mitch Campbell, Regional Wildlife Biologist, DoE, Arviat, NU.
- Project design and implementation.
- Jonathon Pameolik, Regional Wildlife Technician, DoE, Arviat, NU.
- Coordinator of project logistics.

Partners:

The proposed study is partnered with the NWMB, and HTOs Nunavut wide.

Literature Cited:

- Pritchard, J.K., M. Stephens, and P. Donnelly. 2000. Inference of population structure using multilocus genotype data. *Genetics* 155:945-959.
- Proctor, M.F. 2003. Genetic Analysis of Movement and Population Fragmentation of Grizzly Bears in Southwestern Canada. Ph.D. Thesis. Department of Biological Sciences, University of Calgary. 147pp.
- Proctor, M.F., and D. Paetkau. ,2004. A Genetic-based Spatial Analysis of Grizzly Bears in Alberta. Alberta Fish and Wildlife Division. 27pp.
- Raymond, M., and F. Rousset. 1995. GENEPOP Version 1.2: population genetics software for exact tests and ecumenicism. *Journal of Heredity* 86:248-249.
- Zittlau, A.K. 2004. Population Genetic Analyses of North American Caribou (*Rangifer tarandus*). Ph.D. Thesis. Department of Biological Sciences, University of Alberta. 187pp.

Budget Details:

1. Shipping has yet to be accrued.
2. Sample shipping to genetics lab has yet to be accrued.
3. DoE covered the costs of half the genetic analysis in March 2006. Current expected expenditures reflect remaining analysis costs in 2007.
4. Sample collection costs were partially covered in 2006 and for 2007 will be covered by an externally funded contaminants program (i.e. genetic samples will be sub sampled from contaminants tissue samples).
5. NWMB carryover of 34,000.00 to 2006/07 fiscal year.

Budget:

Nunavut Caribou and MuskoX Genetics 2006-2007
Project Name: Nunavut Caribou and MuskoX Genetics
Project Code: 117

Year: 2006-2007

	ORIGINAL TOTAL
Travel and Accommodations	
scheduled air	
meals/accom. (north)	
Materials and Supplies	
Purchased Services	
Shipping	
Sample shipping (HTOs) (26 communities X \$50/Community)	\$1,000.00
Sample Shipping (Genetics Lab)	\$500.00
Contract Services	
technical services (south) (Profiling fees and reporting 500 samples X \$100/sample)	\$50,000.00(Committed)
printing/graphics (Toner cartridges + paper/newsletter)	\$500.00
PROJECT TOTAL	\$52,000.00