

ຈ້ະອັດໂຈ້ລຳຕົວຕາວະີຢ໌ ລອດກະປະລ Munarhiliqiyikkut Inuuhiriknirmullu Department of Health and Social Services Ministère de la Santé et des Services sociaux

ROUTINE IMMUNIZATION & TB SKIN TESTING SCHEDULE 2010 Infant and Childhood Series

Age at Vaccination	BCG	НВ	DTaP-IPV-Hib	Pneu- C ¹ (7,10 or 13)	MMR	Var	Pneu-P	Men-C	Tdap	TST	HPV	Flu ²
Birth	х	х										
1 month		X										
2 month			х	х								
4 month			X	Х								
6 month			х	х								
9 month		Х										
12 month					х			х				
15 month				X		Х						
18 month			х		Х							
2-3 years							X					
4-6 years			X (without Hib)							х		
≥9 year (Grade 6)										x	X (1 st visit)	
											X (2 months after 1 st dose)	
											X (6months after 1 st dose)	
14-16 years (Grade 9)								CU ³	x	х		
≥17years (Grade 12)					CU⁴							

BCG	Bacillus Calmette Guerin.	Pneu-P	Pneumococcal polysaccharide. This is a 23- valent				
HB	Hepatitis B.		vaccine.				
DTaP-IPV-Hib	Diphtheria, acellular Pertussis, Tetanus, Polio,	Men-C	Meningococcal conjugate.				
	(given as 'one needle' with Hib).	Tdap	Tetanus, diphtheria, acellular Pertussis.				
Hib	Haemophilus influenza type b.	TST	Tuberculin skin test (not a vaccine).				
Pneu- C1	Pneumococcal conjugate vaccine. There are currently 3 types: 7, 10 & 13-valent	HPV	Human Papillomavirus.				
MMR	Measles, Mumps, Rubella.	Flu	Influenza Vaccine.				
Var	Varicella.	CU ^{3 & 4}	Catch up.				

1. The Pneumococcal conjugate vaccines (Pneu-C) are not interchangeable / substitutable. You are strongly advised to consult your Regional Communicable Disease Coordinators (RCDC).

2. Annual Influenza vaccine may be given after 6 months of age. Previously unvaccinated children <9 years of age require 2 doses of the split virus vaccine, with an interval of 4 weeks. The 2nd dose is not needed if the child has received one or more doses of vaccine during a previous influenza season.

3. Catch up (CU) of Men-C will be given if the child did not previously receive the vaccine. This practice will continue until 2020 when children born after 2006 and above the age of 1 year should all have received this vaccine.

4. Catch up (CU) of MMR will be given if the child did not previously receive 2 doses of the vaccine. This practice will continue until 2015 when all those who received 2 doses of MMR vaccine after 1995 would have entered grade 12.

Multiple injections should be given at different sites. If more than 2 injections are required, 2 injections may be administered into the same muscle separated by at least 2.5cm (see pg 41 of Canadian Immunization guide (CIG), 7th edition).

Administering acetaminophen with vaccination is not routinely recommended.

Consult the CIG, product monographs or your RCDC if you have questions.

Approved for use in Nunavut by Dr. Geraldine Osborne